

Testing Standard and Degenerate Big Bang Nucleosynthesis with BOOMERanG and MAXIMA-1

S. Esposito¹, G. Mangano¹, A. Melchiorri², G. Miele¹, O. Pisanti¹

¹ *Dipartimento di Scienze Fisiche, Università “Federico II”, Napoli, and INFN Sezione di Napoli, Italy.*

² *Dipartimento di Fisica, Università “La Sapienza”, Roma, Italy.*

We test different Big Bang Nucleosynthesis scenarios using the recent results on the Cosmic Microwave Background Anisotropies provided by the BOOMERanG and MAXIMA-1 experiments versus the observed abundances of ${}^4\text{He}$, D and ${}^7\text{Li}$. The likelihood analysis, based on Bayesian approach, shows that, in the case of high deuterium abundance, $Y_D = (2.0 \pm 0.5)10^{-4}$, both standard and degenerate BBN are inconsistent with the CMBR measurement at more than 3σ . Assuming low deuterium abundance, $Y_D = (3.4 \pm 0.3)10^{-5}$, the standard BBN model is still inconsistent with present observations at 2σ level, while the degenerate BBN results to be compatible. Unless systematics effects will be found in nuclide abundances and/or in CMBR data analysis this result may be a signal in favour of new physics like a large chemical potential of the relic neutrino-antineutrino background.

One of the main goals of modern cosmology is the knowledge of the energy density content of the universe. The four parameters Ω_B , Ω_{CDM} , Ω_ν and Ω_Λ , giving, respectively, the baryon, cold dark matter, neutrino and cosmological constant contributions to the total energy density, in unit of the critical density, and the Hubble constant, $H_0 = 100h \text{ Km s}^{-1} \text{ Mpc}^{-1}$, enter as crucial parameters in several cosmological observables. A well known example is provided by the baryon density parameter which plays an essential role in determining the abundances of light nuclides produced in the early universe. An increasing interest has been also devoted to other issues, as structure formation and the anisotropy of the Cosmic Microwave Background Radiation (CMBR). As for our theoretical understanding of the Big Bang Nucleosynthesis (BBN), to test the theoretical models which describe these aspects of the hot Big Bang model, it is essential to have precise measurements of the several $\Omega_i h^2$. Furthermore, combining different cosmological observables, and comparing the way they are able to constrain the $\Omega_i h^2$, allows for a check of the consistency of our present understanding of the evolution of the universe. This can provide new hints on phenomena which took place at the macroscopic cosmological level, or rather related with the very microscopic structure of fundamental interactions.

This letter represents a contribution in this direction. In particular, we do perform a combined analysis of the dependence on the energy fractions $\Omega_B h^2$ and $\Omega_\nu h^2 = N_\nu \Omega_\nu^0 h^2$ for *massless* neutrinos (N_ν standing for the effective neutrino number and $\Omega_\nu^0 h^2$ for the energy contribution of a single $\nu - \bar{\nu}$ specie) of CMBR anisotropies and BBN. This is aimed to test the standard and degenerate BBN scenario, using the recent results of the BOOMERanG [1] and MAXIMA-1 [2] CMBR experiments and the measurements of ${}^4\text{He}$, D and ${}^7\text{Li}$ primordial abundances.

The theoretical tools necessary to achieve this goal

are nowadays rather robust and the accuracy in the predictions of the Big Bang model is remarkably improved. In fact, recently, a new generation of BBN codes have been developed [3–6], which give the ${}^4\text{He}$ mass fraction Y_p with a theoretical error of few per mille. This effort is justified in view of the small statistical error which is now quoted in the Y_p measurements. On the other hand, the theoretical predictions on the CMBR anisotropies angular power spectrum, in the case of primordial *passive* and *coherent* perturbations, expected in the *most general* inflationary model, have also recently reached a 1% level accuracy, with a refined treatment of H, He I, and He II recombination [7].

Concerning the CMBR experimental data, after the COBE satellite first detection of CMBR anisotropies, at scales larger than 5° , and more than 20 independent detections at different frequencies and scales, see e.g. [8], an important new insight is represented by the recent results obtained by the BOOMERanG Collaboration [1]. For the first time, in fact, multifrequency maps of the microwave background anisotropies were realized over a significant part of the sky, with $\sim 10'$ resolution and high signal to noise ratio. The anisotropy power spectrum, C_ℓ , was measured in a wide range of angular scales from multipole $\ell \sim 50$ up to $\ell \sim 600$, with error bars of the order of 10%, showing a peak at $\ell_{peak} = (197 \pm 6)$ with an amplitude $DT_{200} = (69 \pm 8) \mu K$. While the presence of such peak, compatible with inflationary scenario, was already suggested by previous measurements [9], the absence of secondary peaks after $\ell \geq 300$ with a flat spectrum with an amplitude of $\sim 40 \mu K$ up to $\ell \sim 625$ was a new and unexpected result. This result obtained then an impressive confirmation by the MAXIMA-1 [2] experiment up to $\ell \sim 800$.

As far as BBN is concerned, in the last few years many results have been also obtained on light element primordial abundances. The ${}^4\text{He}$ mass fraction Y_p ,

has been measured with a 0.1% precision in two independent surveys, from regression to zero metallicity in Blue Compact Galaxies, giving a *low* value $Y_p^{(l)} = 0.234 \pm 0.003$ [10], and a high one $Y_p^{(h)} = 0.244 \pm 0.002$ [11], which are compatible at 2σ level only, may be due to large systematic errors. As in [4], in our analysis we adopt the more conservative value $Y_p = 0.238 \pm 0.005$. A similar controversy holds in D measurements, where observations in different Quasars Absorption line Systems (QAS) lead to the incompatible results $Y_D^{(l)} = (3.4 \pm 0.3) 10^{-5}$ [12], and $Y_D^{(h)} = (2.0 \pm 0.5) 10^{-4}$ [13]. We will perform our analysis for both low and high D data. Finally, the most recent estimate for ${}^7\text{Li}$ primordial abundance, from the *Spite plateau* observed in the halo of POP II stars, gives $Y_{Li} = (1.73 \pm 0.21) 10^{-10}$ [14]. The light nuclide yields strongly depend on the baryon matter content of the universe, $\Omega_B h^2$. High values for this parameter result in a larger ${}^4\text{He}$ mass fraction and a lower deuterium number density. In particular, assuming a standard BBN scenario, i.e. vanishing neutrino chemical potentials, the likelihood analysis gives, at 95% C.L. [4],

$$\begin{aligned} \text{low } D \quad \Omega_B h^2 &= .017 \pm .003 \quad 1.7 \leq N_\nu \leq 3.3 \quad , \\ \text{high } D \quad \Omega_B h^2 &= .007^{+0.007}_{-0.002} \quad 2.3 \leq N_\nu \leq 4.4 \quad . \end{aligned} \quad (1)$$

As already pointed out by several authors [15–17], these values for $\Omega_B h^2$, though in the correct order of magnitude, are however somehow smaller than the baryon fraction which more easily fit the CMBR data. In fact, as mentioned, while the narrow first peak around $l \sim 200$ is a confirmation of the inflationary paradigm, a flat universe with adiabatic perturbations, the lack of observation of a secondary peak at smaller scales raises new intriguing questions about the values of the cosmological parameters. In particular, this result may be a signal in favour of a larger $\Omega_B h^2 \sim 0.03$, since increasing the baryon fraction enhances the odd peaks only. In this respect the measurement of the third peak at larger multipole moments is extremely important. Though it is fair to say that a further analysis of the BOOMERanG and MAXIMA-1 data, as well as new data from future experiment, are needed to clarify this issue, it is however timely to study how our theoretical understanding of BBN can be reconciled with the larger values of $\Omega_B h^2$ suggested by CMBR data.

In figure 1 we start addressing the problem quantitatively, by reporting our theoretical predictions on the several abundances, together with the corresponding experimental values, and the CMBR 68% C.L. bound on $\Omega_B h^2 = 0.030^{+0.007}_{-0.004}$, obtained by our joint analysis of both BOOMERanG and MAXIMA-1 data (see below for the details). The BBN predictions are obtained in the standard scenario for $N_\nu = 3$. Unless systematics effects will be found in nuclide abundances and/or in CMBR data analysis, we see that there is a rather sensible disagreement between the values of the baryon energy density required by standard BBN




FIG. 1. The theoretical abundances for D , ${}^3\text{He}$, ${}^4\text{He}$ and ${}^7\text{Li}$ nuclides, for $N_\nu = 3$, versus the experimental results, represented by the horizontal bands, are reported. The long vertical band represents the 68% C.L. interval for $\Omega_B h^2$ obtained by the BOOMERanG and MAXIMA-1 data analysis.

and by CMBR data. In particular, from the 68% C.L. CMBR result for $\Omega_B h^2$, we can derive the following constraints on the Y_i ,

$$\begin{aligned} 0.2487 &< Y_{4\text{He}} < 0.2536 \\ 9.23 \cdot 10^{-6} &< Y_D < 2.32 \cdot 10^{-5} \\ 7.92 \cdot 10^{-6} &< Y_{3\text{He}} < 9.85 \cdot 10^{-6} \\ 5.45 \cdot 10^{-10} &< Y_{Li} < 1.25 \cdot 10^{-9} \quad , \end{aligned} \quad (2)$$

which are systematically different from the available measured values. We stress once again that our theoretical analysis is accurate, at worst, at the level of few percent*.

It has already been stressed [17,4] that a simple way to improve the agreement of observed nuclide abundances with $\Omega_B h^2 \geq 0.02$ is to assume non vanishing neutrino chemical potentials at the BBN epoch, a

*It is important to note that this result is obtained following a Bayesian approach in the analysis. Assuming the standard BBN, the best fit model for the CMBR spectrum has a normalized $\chi^2 \sim 50$, over ~ 40 degrees of freedom. Being the χ^2 slightly greater than one, a CMB analysis using the BBN prior as in [15] is still perfectly consistent.

scenario already extensively studied in the past [18]. The effect of neutrino chemical potentials μ_α , with α the neutrino specie, is twofold. A non-vanishing $\xi_\alpha = \mu_{\nu_\alpha}/T_\nu$, contribute to N_ν as

$$N_\nu = 3 + \Sigma_\alpha \left[\frac{30}{7} \left(\frac{\xi_\alpha}{\pi} \right)^2 + \frac{15}{7} \left(\frac{\xi_\alpha}{\pi} \right)^4 \right] \quad , \quad (3)$$

implying a larger expansion rate of the universe with respect to the non-degenerate scenario, and a higher value for the neutron to proton density ratio at the freeze-out. Furthermore, a positive (negative) value for ξ_e means a larger (smaller) number of ν_e with respect to $\bar{\nu}_e$, thus enhancing (lowering) $n \rightarrow p$ processes. Notice that extra relativistic degrees of freedom, like light sterile neutrinos, would contribute to N_ν as well, and in this respect BBN cannot distinguish between their contribution to the total universe expansion rate and the one due to neutrino degeneracy. Therefore our estimates for N_ν can only represent an upper bound for the total neutrino chemical potentials. However, it is also worth stressing that a large $N_\nu > 3$, *does* indeed require a positive ξ_e , since to end up with the observed values for the Y_i , a fine tuned balance between ξ_e and N_ν is required [18,4].

Increasing N_ν also weakly affects the CMBR anisotropy spectrum in two ways. The growth of perturbations inside the horizon is in fact lowered, resulting in a decay of the gravitational potential and hence in an increase of the anisotropy near the first peak. Moreover, the size of horizon and sound horizon at the last scattering surface is changed, and this, with additional effects in the damping, varies the amplitude and position of the other peaks, see e.g. [19].

To test the degenerate BBN scenario we have performed a likelihood analysis of the data. First, to constrain the values of the parameter set $(\xi_e, N_\nu, \Omega_B h^2)$ from the data on ${}^4\text{He}$, D and ${}^7\text{Li}$ we define a *total likelihood function*,

$$\mathcal{L}_{Nucl}(N_\nu, \Omega_B h^2, \xi_e) = L_D L_{{}^4\text{He}} L_{{}^7\text{Li}} \quad , \quad (4)$$

as described in Ref. [4]. For a fully degenerate BBN, since the effect of a positive ξ_e can be compensated by larger N_ν , $\mathcal{L}_{Nucl}(N_\nu, \Omega_B h^2, \xi_e)$ may sensibly differ from zero in a region with rather large values of N_ν . We have chosen to constrain this parameter to be $N_\nu < 16$. This upper limit has been considered after checking that it is well outside the 95% upper limit on N_ν from the BOOMERanG and MAXIMA-1 data (again, see below). The other two parameters are chosen in the following ranges, $-1 \leq \xi_e \leq 1$ and $0.004 \leq \Omega_B h^2 \leq 0.110$.

Since CMBR spectrum is not sensible to ξ_e alone, we have marginalized over ξ_e . All nuclide abundances have been evaluated using the new BBN code described in [4], while the theoretical uncertainties $\sigma_i^{th2}(N_\nu, \Omega_B h^2)$ are found by linear propagation of the errors affecting the various nuclear rates entering in the nucleosynthesis reaction network [6].

The CMBR data analysis methods have been already extensively described in various papers [15]. The anisotropy power spectrum, was estimated in 12 orthogonalized (independent) bins between $\ell = 50$ and $\ell = 650$ by the BOOMERanG experiment and in 10 bins, from $\ell = 70$ to $\ell = 750$, by the MAXIMA-1 experiment. The likelihood function of the CMBR signal C_B inside the bins is well approximated by an offset lognormal distribution, such that the quantity $D_B = \ln(C_B + x_B)$ (where x_B is the offset correction) is a gaussian variable [20]. The likelihood for a given cosmological model is then defined by $-2\ln L = (D_B^{th} - D_B^{ex}) M_{BB'} (D_B^{th} - D_B^{ex})$, where $M_{BB'}$ is a matrix that defines the noise correlations and D_B^{th} is the offset lognormal theoretical band power. The theoretical D_B^{th} were generated using fast and accurate Boltzmann solvers [21]. Our database of models is sampled in *physical variables* as in [15], but we only consider flat models, $h = 0.65 \pm 0.2$ and the effective number of neutrinos is allowed to vary up to $N_\nu = 20$. Following [22] we can therefore constrain the parameter of interest, $\Omega_B h^2$ and N_ν , by finding the remaining “nuisance” parameters which maximize them. As mentioned, at 95 and 99% C.L. we found $N_\nu \leq 13$ and $N_\nu \leq 16$, respectively. A similar bound, using BOOMERanG data only, has been obtained in [23].

In figure 2 we have summarized the second main result of our analysis. In the $\Omega_B h^2 - N_\nu$ plane we show the 95% C.L. likelihood regions for both the high and low D measurements, as well as the analogous contours for standard BBN, obtained running our code with $\xi_e = 0$. In the same plot we show the 68 and 95 % C.L. regions obtained by CMBR data.

As a first comment, the standard BBN, $\xi_e = 0$, and CMBR data analysis lead to quite different values for $\Omega_B h^2$. This can be clearly seen from the reported 95% results, but we have verified that the 99% C.L. contour for high D has no overlap with the region picked up by BOOMERanG and MAXIMA-1 data, and a very marginal one for low D . For the degenerate scenario, increasing N_ν , the allowed intervals for $\Omega_B h^2$ shift towards larger values. However the high D values require a baryon content of the universe energy density which is still too low, $\Omega_B h^2 \leq 0.018$, to be in agreement with CMBR results. A large overlap is instead obtained for the low D case, whose preferred $\Omega_B h^2$ span the range $0.012 \leq \Omega_B h^2 \leq 0.036$. As expected, a larger N_ν helps in improving the agreement with the high CMBR $\Omega_B h^2$ value, but is important to stress that a large value for N_ν is not preferred by the CMBR data alone, being, in this case, the best fit $N_\nu \sim 3$. If we only consider the 95% overlap region we get the following conservative bounds:

$$4 \leq N_\nu \leq 13 \quad , \quad 0.024 \leq \Omega_B h^2 \leq 0.034 \quad . \quad (5)$$

In this region ξ_e varies in the range $0.07 \leq \xi_e \leq 0.43$. As we said, values $N_\nu \geq 3$, as suggested from our analysis, can be either due to weak interacting neutrino degeneracy, or rather to other unknown relativistic




FIG. 2. The 95% C.L. contours in the $\Omega_B h^2 - N_\nu$ plane compatible with degenerate BBN (large bands) and standard BBN (small regions) are plotted for both high D (left) and low D (right). The same contours for 68% and 95% C.L. from BOOMERanG and MAXIMA-1 CMBR data are also reported.

degrees of freedom.

In conclusion, we have shown how a precision analysis of BBN and CMBR data is able to tell us about possible new features of the cosmological model describing the evolution of the universe. We have quantitatively discussed how larger values for the baryonic matter content $\Omega_B h^2$ may be reconciled with BBN predictions in a degenerate scenario. In this respect the observed low value of deuterium more easily fits with the constraints given by BOOMERanG and MAXIMA-1 data. The crucial aspect of our analysis is that this agreement is realized with an effective neutrino degrees of freedom larger than 4 at 95%. This represents an indication in favour of a degenerate neutrino background and/or new particle species contributing to relativistic matter in the universe. As final remark, we should note that our CMB analysis was restricted to a specific class of models with a limited numbers of parameters. Including curvature, a gravity waves background, or removing our priors on h would not move the peak of our likelihood on $\Omega_B h^2$, but would enlarge our C.L. more on the high value side [15]. Including drastical deviations from the standard model like topological defects [24], decoherence in the primordial fluctuations [25] or assuming a less general inflationary model [26] would drastically change the conclusions of our work. Each of these effect leaves a characteristic imprint on CMB, so hopefully with new data available in the near future, as well as further analysis of the BOOMERanG and MAXIMA results, it will be possible to severely scrutinize this result.

One of the author, A. Melchiorri would like to thank P. de Bernardis, P. Ferreira, J. Silk and N. Vittorio for valuable comments and discussions; G. Miele would like to thank CERN TH-Division for support.

- [1] P. de Bernardis et al. (Boomerang Coll.), *Nature* **404** 955 (2000).
- [2] S. Hanany et al. (Maxima Coll.), *Astrophys. J.*, *submitted*, (2000).
- [3] S. Esposito, G. Mangano, G. Miele, and O. Pisanti, *Nucl. Phys.* **B540** 3 (1999); S. Esposito, G. Mangano, G. Miele, and O. Pisanti, *Nucl. Phys.* **B568** 421 (2000).
- [4] S. Esposito, G. Mangano, G. Miele, and O. Pisanti, astro-ph/0005571.
- [5] R.E. Lopez and M.S. Turner, *Phys. Rev.* **D59** 103502 (1999).
- [6] E. Lisi, S. Sarkar, and F.L. Villante, *Phys. Rev.* **D59** 123520 (1999).
- [7] S. Seager, D.D. Sasselov, and D. Scott, *Astrophys. J.* **523** 1 (1999).
- [8] E. Pierpaoli, D. Scott, and M. White, *Science* 2171 (2000); L. Knox and L. Page, *Phys. Rev. Lett.* *in press*, (2000).
- [9] P. D. Mauskopf et al, *Astrophys. J.*, **L59** 536, (2000); A. D. Miller et al., *Astrophys. J.*, **L1** 524, (1999).
- [10] K.A. Olive and G. Steigman, *Astrophys. J. Suppl.*, **97** 49 (1995), and references therein.
- [11] Y.I. Izotov and T.X. Thuan, *Astrophys. J.*, **500** 188 (1998).
- [12] S. Burles and D. Tytler, *Astrophys. J.*, **499** 689 (1998).
- [13] For the most recent analysis see J.K. Webb, R.F. Carswell, K.M. Lanzetta, R. Ferlet, M. Lemoine, A. Vidal-Madjar, and D.V. Bowen, *Nature*, **388** 250 (1997).
- [14] P. Bonifacio and P. Molaro, *Mon. Not. Roy. Astron. Soc.*, **285** 847 (1997).
- [15] A.E. Lange et al. (Boomerang Coll.), astro-ph/0005004; M. Tegmark and M. Zaldarriaga, astro-ph/0004393, A. H. Jaffe et al., astro-ph/0007333; W. Kinney, A. Melchiorri, A. Riotto, astro-ph/0007375.
- [16] M. Le Dour, M. Douspis, J.G. Barlett, A. Blanchard, astro-ph/0004283; M.White, D. Scott, and E. Pierpaoli, astro-ph/0004385.
- [17] J. Lesgourgues and S. Pastor, *Phys. Rev.*, **D60** 103521 (1999); W.H. Kinney and A. Riotto, *Phys. Rev. Lett* **83** 3366 (1999); J. Lesgourgues and M. Peloso, astro-ph/0004412; M. Orito, T. Kajino, G.J.Mathews, and R.N.Boyd, astro-ph/0005446.
- [18] H. Kang and G. Steigman, *Nucl. Phys.*, **B372** 494 (1992).
- [19] W. Hu et al., *Phys. Rev.*, **D52** 5498 (1995).
- [20] J.R. Bond, A. H. Jaffe, and L. Knox, *Astrophys. J.*, **533** 19 (2000).
- [21] U. Seljak and M. Zaldarriaga, *Astrophys. J.*, **469** 437 (1996); A. Lewis, A. Challinor, and A. Lasenby, *Astrophys. J.*, *in press* (2000).
- [22] L. Knox and S. Dodelson, *Phys. Rev. Lett.*, **84** 3523 (2000); A. Melchiorri et al., *Astrophys. J.*, **536** L63 (2000).
- [23] S. Hannestad, astro-ph/0005018.
- [24] F. R. Bouchet et al, astro-ph/0005022.
- [25] A. Lewin, and A. Albrecht, astro-ph/9908061.
- [26] T. Kanazawa et al, astro-ph/0006445.